




NOAA Technical Memorandum

NMFS-SEFC-215

Length-Weight Relationships of Selected Marine Reef Fishes
from the Southeastern United States and the Caribbean



DECEMBER 1988

James A. Bohnsack
Douglas E. Harper

U. S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Marine Fisheries Service
Southeast Fisheries Center
Miami, Florida 33149



NOAA Technical Memorandum

NMFS-SEFC-215

Length-Weight Relationships of Selected Marine Reef Fishes
from the Southeastern United States and the Caribbean

BY

James A. Bohnsack and Douglas E. Harper

U. S. DEPARTMENT OF COMMERCE
C. William Verity, Jr., Secretary

National Oceanic and Atmospheric Administration
William E. Evans, Administrator

National Marine Fisheries Service
James W. Brennan, Assistant Administrator for Fisheries

DECEMBER 1988

This Technical Memorandum series is used for documentation and timely communication of preliminary results, interim reports, or similar special-purpose information. Although the memoranda are not subject to complete formal review, editorial control, or detailed editing, they are expected to reflect sound professional work.

NOTICE

The National Marine Fisheries Service (NMFS) does not approve, recommend or endorse any proprietary product or material mentioned in this publication. No reference shall be made to NMFS, or to this publication furnished by NMFS, in any advertising or sales promotion which would indicate or imply that NMFS approves, recommends, or endorses any proprietary product or proprietary material mentioned herein, or which has as its purpose an intent to cause directly or indirectly the advertised product to be used or purchased because of this NMFS publication.

This report should be cited as follows:

Bohnsack, J.A. and D.E. Harper. 1988. Length-weight relationships of selected marine reef fishes from the southeastern United States and the Caribbean. NOAA Technical Memorandum NMFS-SEFC-215, 31 p.

Copies may be obtained by writing:

National Technical Information Service
5248 Port Royal Rd.
Springfield, VA 22161

TABLE OF CONTENTS

Abstract	1
Introduction	1
Purpose	1
Methods	1
Results	2
Discussion	2
Acknowledgments	2
Literature Cited	3
Table 1	4 - 7
Table 2	8 - 9
Table 3	10 - 14
Appendix A	15 - 31

Length-Weight Relationships of Selected Marine Reef Fishes from the Southeastern United States and the Caribbean

by James A. Bohnsack and Douglas E. Harper

*National Oceanic and Atmospheric Administration
National Marine Fisheries Service
Southeast Fisheries Center
Miami Laboratory
75 Virginia Beach Drive
Miami, Florida 33149*

restricted to single or a few commercial species (e.g., Starck, 1970; Campbell, 1984).

Length-to-weight conversion formulae would be helpful in estimating reef fish biomass. Biomass data are important for studying and modeling ecosystem structure, trophic relationships, population dynamics, species importance, stock characteristics, and fisheries exploitation. Lengths converted to weights can be used to estimate catch-at-age for fisheries analysis. In addition, conversion formulae may help law enforcement personnel estimate total catch weights for legal purposes.

ABSTRACT

Formulae are provided for converting length estimates to wet weight biomass for 178 fish species from the southeast coast of the U.S., Gulf of Mexico, and the Caribbean. The formulae were empirically derived from measurements of 31,886 fishes. Species observed but lacking length-weight data are identified. Data contributions are solicited for future updates.

PURPOSE

Here we provide a preliminary list of empirically derived length-to-weight relationships based on data collected during fishery research in the Caribbean and off the southeastern United States. Our intent is to provide a convenient reference of length-weight formulae that can be used to convert size-frequency observations to biomass estimates for comprehensive fish community and stock comparisons.

INTRODUCTION

A wide variety of visual censusing methods have been developed for non-destructive fishery-independent stock assessments. Most visual surveys of reef fishes have been limited to listing species or counting individuals (e.g., Brock, 1954; Jones and Thompson, 1978; Kimmel, 1985). A few visual census methods provide length estimates either for selected species (Bell, et al., 1985) or for general fish community structure (Bohnsack and Bannerot, 1986). Although fish length can be estimated underwater, use of these data have been limited because of a lack of information on relationships between length and weight for a majority of species. The mathematics of length-weight relationships has been reviewed by Weatherley and Gil (1987). Published data usually have been

METHODS

Scientists from the Miami Laboratory, SEFC, have opportunistically collected biological data on reef fishes from southern Florida from 1980 to the present. All fish were measured with a standard fish measuring board graduated in millimeters (mm). Fork lengths (FL) were recorded to the nearest mm by measuring from the tip of the snout with the mouth closed, to the end of the middle-most caudal ray. Total length (TL), measured to the longest caudal ray, was used for species with rounded or truncated caudal fins. Weights were measured to the nearest 0.1 g on a laboratory triple beam balance whenever the fish could be brought into the laboratory. Fish processed in the field were weighed on spring-tension cylindrical scales to the nearest 10 g.

In addition to southern Florida fishes, length and weight information for reef fish were obtained from the 1985 Caribbean Biostatistical Database. Bohnsack et. al. (1986) provides detailed analysis and information concerning this database. For the purposes of this paper, records for all reef fish species were extracted and used in the regression computations. Outliers, data outside of the general length-weight trend area, were manually omitted from the analysis.

Weight-length relationships were calculated by fitting a regression line to the equation:

$$\log W = \log a + b \log L$$

which is equivalent to the equation:

$$W = aL^b$$

where W is weight in grams, and L is length in millimeters and a and b are constants. The regression constants, standard error of the constants, and coefficient of determination (R^2) were obtained using the least squares data regression function of the LOTUS 1-2-3 software.

Regressions for species with less than four actual measurements were derived by forcing the intercept through a point close to the origin by adding a "zero" data point.

RESULTS

Regression formulae are provided for 174 fish species representing a total of 7,503 individuals from the South Atlantic/ Gulf of Mexico (southern Florida) (Table 1) and 32 fish species representing a total of 24,383 individuals from the Caribbean (Table 2). Graphical representations of size-distribution and length-weight relationships for 50 southern Florida species with more than 30 measurements are presented in Appendix A so that users can assess the range and precision of the data. Formulae for species with fewer measurements should be used cautiously. Species observed in visual reef fish samples but lacking adequate length- to-weight data are identified in Table 3.

DISCUSSION

Data include most, but not all, species that have been observed during visual censuses of reef fish

habitats by the staff at the Miami Laboratory. Many species are transients or visiting species and are not considered reef fish by traditional criteria (i.e., Starck, 1968). These species are included, however, because they potentially can be major influences on reef fish community structure.

Biomass is considered an important variable in ecological studies. Numbers of fishes used alone tend to give excessive importance to small species and juvenile fishes. Biomass may be more representative of community structure although it may give excessive importance to very large individuals. Numbers, biomass, and frequency-of-occurrence can be combined to provide an "importance value" for particular species in ecological studies (Brower and Zar, 1977).

We anticipate that the formulae and length-weight data will be useful, but will also indicate gaps in our knowledge. We anticipate publishing more comprehensive periodic updates. We encourage and solicit contributions of data for this effort, particularly for species with few data or lacking data. All contributions will be acknowledged. If sufficient interest exists, we may expand future editions to include other fishery parameters, various length-to-length relationships, regional length-to-weight relationships, and summaries of published length to weight relationships. Suggestions are welcome.

ACKNOWLEDGMENTS

We thank the following people in alphabetical order for providing raw data: Christopher Holt, Ken Lindeman, and David McClellan. Organizations that contributed to this effort include the Newfound Harbor Marine Institute, Big Pine Key, Florida, the University of Miami, Miami, Florida, the Miami Seaquarium, Miami, Florida, and the Caribbean Biostatistical Sampling Program.

LITERATURE CITED

- BELL, J.D., G.J.S. CRAIK, D.A. POLLARD, and B.C. RUSSELL
1985. Estimating length frequency distributions of large reef fish underwater. *Coral Reefs* 4:41-44.
- BOHNSACK, J.A. and S.P. BANNEROT.
1986. A stationary visual census technique for quantitatively assessing community structure of coral reef fishes. *NOAA Tech. Rep. NMFS* 41. 15 p.
- BOHNSACK, J.A. , D.L. SUTHERLAND, A. BROWN, D.E. HARPER, and D.B. McCLELLAN.
1986. An analysis of the Caribbean biostatistical database for 1985. *Coastal Resources Division Report*, Contribution No. CRD-86/87-10.
- BROCK, V.E.
1954. A preliminary report on a method of estimating reef fish populations. *J. Wildl. Mgmt.* 18:297-308.
- BROWER, J.E. and J.H. ZAR.
1977. Field and laboratory methods for general ecology. Wm. C. Brown Co., Publishers. Dubuque, Iowa, 194 p.
- CAMPBELL, P.
1984. Weight-total length and length-length relationships for four saltwater fishes. Management Data Series No. 62. Texas Parks and Wildlife Department. Austin, Texas. 7 p.
- JONES, R.S. and M.J. THOMPSON.
1978. Comparison of Florida reef fish assemblages using a rapid visual technique. *Bull. Mar. Sci.* 28:159-172.
- KIMMEL, J.J.
1985. A new species-time method for visual assessment of fishes and its comparison with established methods. *Env. Biol. Bish.* 12:23-32.
- ROBINS, C.R., R.M. BAILEY, C.E. BOND, J.R. BROOKER, E.A. LACHNER, R.N. LEA, and W.B. SCOTT.
1980. A list of common and scientific names of fishes from the United States and Canada. Fourth edition. American Fisheries Society Special Publication No. 12. 174 p.
- STARCK, W.A., II.
1968. A list of fishes of Alligator Reef, Florida with comments on the nature of the Florida reef fish fauna. *Undersea Biol.* 1:5-36.
- STARCK, W.A., II.
1970. Biology of the gray snapper, *Lutjanus griseus* (Linneaus), in the Florida Keys. *Stud. Trop. Oceanogr.* Miami 10:11-150.
- WEATHERLY, A.H. and H.S. GILL.
1987. The biology of fish growth. Academic Press, Harcourt Brace Jovanovich, Publishers. New York. 443 p.
- STARCK, W.A., II.
1968. A list of fishes of Alligator Reef, Florida with

Table 1. - Summary of southern Florida reef fish weight-regressions for 7,503 fish.

Regression formula: $\log \text{WEIGHT(gms)} = \log a + b \log \text{LENGTH(mm)}$.

Names of fishes are from Robins et. al. (1980).

* - Species with less than four actual fish measurements. The regression values presented can be only considered to represent very crude estimates. The regression lines were derived by forcing the intercept through a point at or close to the origin by adding a "zero" data point.

SPECIES	COMMON NAME	NO.	TYPE	SIZE (mm)			log a	b	R^2	STANDARD DEVIATION	
				MIN.	MAX.					b	log a
1 <u>Abudefduf saxatilis</u>	Sergeant major	35	FL	18	143	-4.7859	3.1420	0.98	0.0864	0.1359	
2 * <u>Acanthemblemaria aspera</u>	Roughhead blenny	1	TL	30	30	-5.0750	2.9625	1.00	0.0000	0.0000	
3 <u>Acanthurus bahianus</u>	Ocean surgeon	29	FL	35	243	-4.6005	2.9752	0.99	0.0666	0.0771	
4 <u>Acanthurus chirurgus</u>	Doctorfish	61	FL	39	304	-5.9255	3.5328	0.94	0.1154	0.1438	
5 <u>Acanthurus coeruleus</u>	Blue tang	394	FL	35	304	-4.2165	2.8346	0.97	0.0261	0.0775	
6 <u>Aluterus schoepfii</u>	Orange filefish	53	TL	277	548	-3.3770	2.3443	0.78	0.1732	0.0671	
7 <u>Aluterus scriptus</u>	Scrawled filefish	71	TL	193	615	-1.8982	1.8136	0.89	0.0781	0.0908	
8 <u>Amblycirrhitus pinos</u>	Redspotted hawkfish	25	FL	275	452	-6.0169	3.4266	0.87	0.2712	0.0610	
9 <u>Anisotremus surinamensis</u>	Black margate	21	FL	247	488	-5.6212	3.3916	0.85	0.3223	0.0968	
10 <u>Anisotremus virginicus</u>	Porkfish	187	FL	17	298	-4.9963	3.1674	0.99	0.0239	0.0816	
11 <u>Apogon maculatus</u>	Flamefish	22	FL	16	65	-4.8752	3.0734	0.97	0.1110	0.0907	
12 <u>Apogon pseudomaculatus</u>	Twospot cardinalfish	49	FL	18	68	-4.6413	2.9434	0.93	0.1156	0.1123	
13 * <u>Archosargus probatocephalus</u>	Sheepshead	1	FL	271	271	-4.8175	3.0912	1.00	0.0000	0.0000	
14 <u>Archosargus rhomboidalis</u>	Sea bream	18	FL	42	254	-4.8471	3.1021	1.00	0.0518	0.0477	
15 * <u>Astrapogon stellatus</u>	Conchfish	1	TL	57	57	-4.8463	3.0769	1.00	0.0000	0.0000	
16 * <u>Aulostomus maculatus</u>	Trumpetfish	1	TL	209	209	-5.2686	2.8657	1.00	0.0000	0.0000	
17 <u>Balistes capricus</u>	Gray triggerfish	233	FL	158	536	-4.5359	2.9352	0.93	0.0549	0.0762	
18 <u>Balistes vetula</u>	Queen triggerfish	22	FL	165	425	-4.5638	2.9903	0.96	0.1388	0.0550	
19 <u>Blennius cristata</u>	Molly miller	16	TL	22	73	-3.9658	2.4144	0.92	0.1873	0.0938	
20 * <u>Bodianus rufus</u>	Spanish hogfish	2	FL	252	331	-4.8936	3.0532	1.00	0.0012	0.0045	
21 <u>Bothus lunatus</u>	Peacock flounder	10	TL	81	469	-5.1974	3.1894	0.99	0.0978	0.0707	
22 <u>Calamus bajonado</u>	Jolthead porgy	130	FL	192	481	-4.1758	2.8175	0.92	0.0730	0.0511	
23 <u>Calamus calamus</u>	Saucereye porgy	207	FL	158	310	-4.1688	2.8009	0.84	0.0850	0.0640	
24 <u>Calamus penna</u>	Sheepshead porgy	28	FL	165	369	-3.7826	2.6663	0.83	0.2401	0.1017	
25 <u>Calamus proridens</u>	Littlehead porgy	47	FL	134	272	-3.5329	2.5299	0.75	0.2206	0.1140	
26 * <u>Callionymus bairdii</u>	Lancer dragonet	1	TL	16	16	-4.7580	3.1210	1.00	0.0000	0.0000	
27 <u>Cantherhines macrocerus</u>	Whitespotted filefish	10	TL	233	395	-3.9037	2.6534	0.90	0.3109	0.0633	
28 <u>Cantherhines pullus</u>	Orangespotted filefish	12	TL	151	200	-3.7282	2.5632	0.91	0.2508	0.0314	
29 <u>Canthidermis sufflamen</u>	Ocean triggerfish	34	FL	363	555	-4.8095	3.0554	0.97	0.1023	0.0343	
30 <u>Caranx bartholomaei</u>	Yellow jack	34	FL	202	705	-4.4953	2.9085	0.99	0.0536	0.0484	
31 <u>Caranx cryos</u>	Blue runner	235	FL	160	458	-3.9699	2.6900	0.84	0.0782	0.1042	
32 <u>Caranx hippos</u>	Crevalle jack	18	FL	260	647	-4.0201	2.7344	0.99	0.0706	0.0422	
33 <u>Caranx ruber</u>	Bar jack	34	FL	32	377	-5.3687	3.2370	0.98	0.0754	0.0800	
34 <u>Centropristes oxyurus</u>	Bank sea bass	10	TL	192	299	-3.6555	2.5990	0.96	0.1944	0.0422	
35 <u>Centropristes striata</u>	Black sea bass	7	TL	215	328	-3.6557	2.4683	0.97	0.1903	0.0350	
36 <u>Chaetodiperus faber</u>	Atlantic spadefish	43	FL	201	396	-3.7175	2.6836	0.91	0.1348	0.0495	
37 <u>Chaetodon capistratus</u>	Foureye butterflyfish	6	TL	43	115	-4.8475	3.1897	1.00	0.0323	0.0122	
38 <u>Chaetodon ocellatus</u>	Spotfin butterflyfish	80	TL	103	181	-4.4820	2.9838	0.96	0.0732	0.0296	
39 <u>Chaetodon sedentarius</u>	Reef butterflyfish	34	TL	37	134	-4.6749	3.0760	0.88	0.1990	0.1031	
40 <u>Chaetodon striatus</u>	Banded butterflyfish	13	TL	57	143	-4.7940	3.1395	0.99	0.1043	0.0599	
41 * <u>Chilomycterus schoepfii</u>	Stripped burrfish	1	FL	212	212	-4.7512	3.1244	1.00	0.0000	0.0000	
42 <u>Coryphopterus glaucofraenum</u>	Bridled goby	18	TL	17	48	-4.8489	2.9674	0.93	0.2033	0.1188	
43 * <u>Cryptotomus roseus</u>	Bluelip parrotfish	3	TL	12	69	-4.4794	3.1816	0.99	0.2328	0.6985	
44 * <u>Dasyatis americana</u>	Southern stingray	2	TL	651	4150	-5.5182	2.6724	0.99	0.2951	1.2681	
45 <u>Diodon holocanthus</u>	Balloonfish	162	TL	112	430	-3.1948	2.3979	0.73	0.1157	0.1126	
46 <u>Diodon hystrix</u>	Porcupinefish	30	TL	103	345	-2.5498	2.2763	0.99	0.0384	0.0348	
47 <u>Diplectrum formosum</u>	Sand perch	170	FL	39	728	-5.0229	3.0781	0.96	0.0463	0.1115	
48 <u>Echeneis naucrates</u>	Sharksucker	5	TL	430	479	-3.0099	2.1128	0.98	0.1651	0.0060	
49 * <u>Elops saurus</u>	Ladyfish	2	FL	293	413	-5.0651	2.9680	1.00	0.0117	0.0434	
50 * <u>Enneanectes pectoralis</u>	Redeye triplefin	3	TL	23	34	-4.8980	3.0502	1.00	0.0158	0.0474	

Table 1. (continued) - Summary of southern Florida reef fish weight-regressions for 7,503 fish.

Regression formula: $\log \text{WEIGHT(gms)} = \log a + b \log \text{LENGTH(mm)}$.

Names of fishes are from Robins et. al. (1980).

* - Species with less than four actual fish measurements. The regression values presented can be only considered to represent very crude estimates. The regression lines were derived by forcing the intercept through a point at or close to the origin by adding a "zero" data point.

SPECIES	COMMON NAME	SIZE (mm)					STANDARD DEVIATION			
		NO.	TYPE	MIN.	MAX.	log a	b	R ²	b	log a
51 * <i>Epinephelus cruentatus</i>	Graysby	2	TL	219	246	-4.9123	3.0439	1.00	0.0030	0.0107
52 <i>Epinephelus guttatus</i>	Red hind	20	TL	139	255	-5.0680	3.1124	0.92	0.2174	0.0778
53 <i>Epinephelus morio</i>	Red grouper	47	TL	82	566	-4.9464	3.0350	0.99	0.0553	0.0585
54 <i>Epinephelus striatus</i>	Nassau grouper	9	TL	203	516	-5.4164	3.2292	0.98	0.1890	0.0810
55 <i>Equetus acuminatus</i>	High-hat	19	TL	28	199	-5.2620	3.2017	1.00	0.0492	0.0534
56 <i>Equetus lanceolatus</i>	Jackknife-fish	8	TL	93	223	-6.8068	3.8444	1.00	0.0851	0.0278
57 * <i>Eucinostomus havana</i>	Bigeye mojarra	1	FL	116	116	-4.8940	3.0530	1.00	0.0000	0.0000
58 <i>Gerres cinereus</i>	Yellowfin mojarra	4	FL	52	257	-4.8195	3.0843	1.00	0.1145	0.0650
59 <i>Ginglymostoma cirratum</i>	Nurse shark	16	TL	229	1070	-4.8681	2.8918	0.93	0.2143	0.1516
60 <i>Gnatholepis thompsoni</i>	Goldspot goby	9	TL	15	52	-6.2207	3.7669	0.96	0.3071	0.1676
61 * <i>Gobiesox strumosus</i>	Skilletfish	1	TL	42	42	-4.9281	3.0359	1.00	0.0000	0.0000
62 <i>Gobiosoma oceanops</i>	Neon goby	19	TL	23	45	-5.2341	3.1370	0.95	0.1823	0.0681
63 <i>Gymnothorax funebris</i>	Green moray	9	TL	597	2134	-5.2443	2.8560	0.94	0.2623	0.1475
64 <i>Gymnothorax moringa</i>	Spotted moray	22	TL	403	886	-6.1561	3.1577	0.90	0.2301	0.0931
65 * <i>Gymnothorax nigromarginatus</i>	Blackedge moray	1	TL	486	486	-5.3194	2.8403	1.00	0.0000	0.0000
66 * <i>Gymnothorax vicinus</i>	Purplemouth moray	1	TL	758	758	-5.2478	2.8761	1.00	0.0000	0.0000
67 <i>Haemulon album</i>	Margate	11	FL	162	595	-4.8186	3.0423	0.99	0.1154	0.0603
68 <i>Haemulon aurolineatum</i>	Tomtate	198	FL	12	260	-5.2081	3.2077	0.99	0.0186	0.0893
69 * <i>Haemulon carbonarium</i>	Caesar grunt	3	FL	32	37	-4.8879	3.0559	1.00	0.0218	0.0668
70 <i>Haemulon chrysargyreum</i>	Smallmouth grunt	8	FL	140	200	-2.5578	2.1567	1.00	0.0466	0.0060
71 <i>Haemulon flavolineatum</i>	French grunt	61	FL	32	289	-5.0428	3.1581	0.99	0.0392	0.0606
72 <i>Haemulon macrostomum</i>	Spanish grunt	6	FL	174	203	-4.6419	3.0295	0.91	0.4687	0.0233
73 <i>Haemulon melanurum</i>	Cottonwick	70	FL	29	277	-4.5993	2.9527	0.98	0.0565	0.0519
74 <i>Haemulon parrai</i>	Sailor's choice	143	FL	41	300	-4.6947	2.9932	0.99	0.0237	0.0542
75 <i>Haemulon plumieri</i>	White grunt	362	FL	20	298	-5.0781	3.1612	0.99	0.0148	0.0924
76 <i>Haemulon sciurus</i>	Bluestriped grunt	136	FL	26	265	-4.7114	2.9996	0.99	0.0215	0.0705
77 * <i>Haemulon striatum</i>	Striped grunt	3	FL	40	180	-4.8564	3.0988	1.00	0.0963	0.3398
78 <i>Halichoeres bivittatus</i>	Slippery dick	21	TL	36	152	-4.8117	2.9391	0.96	0.1441	0.1389
79 <i>Halichoeres garnoti</i>	Yellowhead wrasse	5	TL	26	105	-5.6591	3.3747	0.99	0.2205	0.1017
80 <i>Halichoeres maculipinna</i>	Clown wrasse	7	TL	15	97	-6.2524	3.6932	0.99	0.1994	0.1414
81 * <i>Halichoeres radiatus</i>	Puddingwife	2	TL	24	36	-4.9221	3.0382	1.00	0.0145	0.0410
82 <i>Hemiramphus brasiliensis</i>	Ballyhoo	4	TL	275	380	-3.3362	2.3555	1.00	0.0717	0.0080
83 <i>Holacanthus bermudensis</i>	Blue angelfish	117	TL	69	427	-4.3950	2.8994	0.97	0.0513	0.0776
84 <i>Holacanthus ciliaris</i>	Queen anglefish	25	TL	29	350	-4.3731	2.9004	1.00	0.0369	0.0559
85 <i>Holacanthus tricolor</i>	Rock beauty	20	TL	42	213	-4.2262	2.8577	0.98	0.0989	0.0925
86 <i>Holocentrus ascensionis</i>	Squirrelfish	108	FL	190	382	-3.6218	2.5596	0.76	0.1403	0.0704
87 * <i>Holocentrus coruscus</i>	Reef squirrelfish	1	FL	51	51	-4.8611	3.0695	1.00	0.0000	0.0000
88 * <i>Holocentrus rufus</i>	Longspine squirrelfish	3	FL	90	175	-4.8825	3.0594	1.00	0.0716	0.2552
89 <i>Hyleurochilus bermudensis</i>	Barred blenny	6	TL	24	47	-4.9426	3.0332	0.95	0.3295	0.0733
90 * <i>Hypoplectrus unicolor</i>	Butter hamlet	4	FL	109	170	-5.1419	3.1818	0.99	0.0346	0.2216
91 * <i>Kyphosus sectatrix</i>	Bermuda chub	2	FL	190	206	-4.8397	3.0801	1.00	0.0034	0.0120
92 <i>Lachnolaimus maximus</i>	Hogfish	228	FL	68	600	-4.6801	2.9880	0.98	0.0318	0.0518
93 <i>Lactophry斯 bicaudalis</i>	Spotted trunkfish	5	TL	136	271	-2.4461	2.0976	0.95	0.2775	0.0644
94 <i>Lactophry斯 polygonia</i>	Honeycomb cowfish	6	TL	200	301	-5.6339	3.3462	0.94	0.4084	0.0553
95 <i>Lactophry斯 quadricornis</i>	Scrawled cowfish	177	TL	88	550	-3.0194	2.2629	0.87	0.0667	0.0785
96 * <i>Lactophry斯 trigonus</i>	Trunkfish	3	TL	292	395	-4.8334	3.0829	1.00	0.0207	0.0814

Table 1. (continued) - Summary of southern Florida reef fish weight-regressions for 7,503 fish.

Regression formula: $\log \text{WEIGHT(gms)} = \log a + b \log \text{LENGTH(mm)}$.

Names of fishes are from Robins et. al. (1980).

* - Species with less than four actual fish measurements. The regression values presented can be only considered to represent very crude estimates. The regression lines were derived by forcing the intercept through a point at or close to the origin by adding a "zero" data point.

SPECIES	COMMON NAME	SIZE (mm)				STANDARD DEVIATION				
		NO.	TYPE	MIN.	MAX.	log a	b	R ²	b	log a
97 <u>Lactophrys triqueter</u>	Smooth trunkfish	117	TL	124	276	-2.7388	2.2302	0.80	0.1042	0.0744
98 <u>Lagodon rhomboides</u>	Pinfish	12	FL	130	212	-5.2350	3.2504	0.97	0.1652	0.0318
99 <u>Lutjanus analis</u>	Mutton snapper	365	FL	116	722	-4.8030	3.0112	0.97	0.0278	0.0579
100 <u>Lutjanus apodus</u>	Schoolmaster	51	FL	38	791	-4.6909	2.9779	0.99	0.0428	0.0516
101 * <u>Lutjanus buccanella</u>	Blackfin snapper	3	FL	30	104	-5.1309	2.9735	0.99	0.2120	0.6800
102 * <u>Lutjanus cyanopterus</u>	Cubera snapper	1	FL	693	693	-4.8799	3.0601	1.00	0.0000	0.0000
103 <u>Lutjanus griseus</u>	Gray snapper	245	FL	53	513	-4.5159	2.8809	0.97	0.0316	0.0537
104 <u>Lutjanus jocu</u>	Dog snapper	5	FL	265	774	-4.3683	2.8574	0.99	0.1567	0.0616
105 <u>Lutjanus mahogoni</u>	Mahogonay snapper	13	FL	220	357	-4.0870	2.7190	0.84	0.3624	0.0761
106 <u>Lutjanus synagris</u>	Lane snapper	254	FL	25	389	-4.3452	2.8146	0.96	0.0351	0.0709
107 <u>Lythrypnus spilus</u>	Bluegold goby	11	TL	16	40	-4.6635	2.7441	0.94	0.2261	0.0808
108 <u>Malacanthus plumieri</u>	Sand tilefish	5	FL	366	427	-4.1981	2.6290	0.93	0.4208	0.0215
109 <u>Malacoctenus macrops</u>	Rosy blenny	5	TL	25	37	-3.8333	2.2229	0.83	0.5814	0.0937
110 <u>Malacoctenus triangulatus</u>	Saddled blenny	4	TL	41	53	-5.3645	3.1883	0.86	0.8935	0.0750
111 <u>Microspathodon chrysurus</u>	Yellowtail damselfish	8	FL	16	58	-4.7033	3.0825	1.00	0.0652	0.0318
112 * <u>Monacanthus ciliatus</u>	Fringed filefish	1	TL	126	126	-4.8264	3.0868	1.00	0.0000	0.0000
113 <u>Monacanthus hispidus</u>	Planehead filefish	115	TL	74	316	-3.9200	2.6178	0.86	0.0991	0.0934
114 <u>Mulloidichthys martinicus</u>	Yellow goatfish	13	FL	223	293	-6.3369	3.6627	0.96	0.2264	0.0310
115 <u>Muraena miliaris</u>	Goldentail moray	6	TL	249	742	-4.5351	2.5736	0.99	0.0916	0.0366
116 <u>Mycteroperca bonaci</u>	Black grouper	21	FL	201	824	-5.3696	3.2051	1.00	0.0390	0.0351
117 <u>Mycteroperca microlepis</u>	Gag	30	FL	419	851	-4.9169	3.0305	0.98	0.0888	0.0382
118 <u>Ocyurus chrysurus</u>	Yellowtail snapper	257	FL	15	363	-4.1108	2.7180	0.96	0.0363	0.0811
119 * <u>Odontoscion dentex</u>	Reef crocker	1	TL	58	58	-4.9854	3.0073	1.00	0.0000	0.0000
120 * <u>Ogcocephalus radiatus</u>	Polka-dot batfish	3	FL	204	255	-4.8749	3.0626	1.00	0.0018	0.0084
121 <u>Ophioblennius atlanticus</u>	Redlip blenny	9	TL	44	82	-3.8680	2.3791	0.87	0.3500	0.0936
122 * <u>Opistognathus whitehursti</u>	Dusky jawfish	1	TL	144	144	-5.0210	2.9895	1.00	0.0000	0.0000
123 <u>Orthopristis chrysoptera</u>	Pigfish	49	FL	156	234	-5.0130	3.1893	0.95	0.1039	0.0237
124 <u>Pagrus pagrus</u>	Red porgy	113	FL	190	393	-4.6955	3.0027	0.95	0.0663	0.0462
125 * <u>Paranthias furcifer</u>	Creole-fish	3	FL	235	295	-4.9130	3.0430	1.00	0.0124	0.0476
126 * <u>Pempheris schomburgki</u>	Glassy sweeper	1	FL	42	42	-4.8557	3.0721	1.00	0.0000	0.0000
127 <u>Phaeoptyx pigmentaria</u>	Dusky cardinalfish	8	FL	25	49	-5.2814	3.2553	0.99	0.0968	0.0274
128 <u>Pomacanthus arcuatus</u>	Gray angelfish	121	TL	16	438	-4.4311	2.9680	0.96	0.0569	0.1309
129 <u>Pomacanthus paru</u>	French angelfish	63	TL	21	413	-4.8182	3.1264	0.99	0.0326	0.0665
130 <u>Pomacentrus fuscus</u>	Dusky damselfish	12	FL	28	79	-4.3479	2.8956	0.89	0.3139	0.1869
131 <u>Pomacentrus leucostictus</u>	Beaugregory	11	FL	30	89	-4.4057	2.8868	0.97	0.1616	0.0742
132 <u>Pomacentrus partitus</u>	Bicolor damselfish	33	FL	14	69	-4.8921	3.1519	0.88	0.2076	0.2271
133 <u>Pomacentrus planifrons</u>	Three spot damselfish	11	FL	14	105	-4.2782	2.8569	0.99	0.1030	0.1075
134 <u>Pomacentrus variabilis</u>	Cocoa damselfish	25	FL	12	74	-4.3258	2.8365	0.96	0.1259	0.1276
135 <u>Priacanthus arenatus</u>	Bigeye	171	FL	199	726	-4.9233	3.0387	0.91	0.0733	0.0665
136 <u>Priacanthus cruentatus</u>	Glasses eye snapper	10	FL	222	345	-1.6587	1.7495	0.72	0.3880	0.0809
137 * <u>Prionotus roseus</u>	Bluespotted searobin	1	TL	142	142	-4.9430	3.0285	1.00	0.0000	0.0000
138 <u>Pseudupeneus maculatus</u>	Spotted goatfish	32	FL	149	290	-4.8231	3.0257	0.91	0.1743	0.0561
139 * <u>Quisquilia hipoliti</u>	Rusty goby	2	TL	36	46	-4.9180	3.0407	1.00	0.0117	0.0344
140 * <u>Rachycentron canadum</u>	Cobia	1	FL	1085	1085	-5.3753	2.8123	1.00	0.0000	0.0000
141 <u>Rhomboplites aurorubens</u>	Vermilion snapper	8	FL	138	245	-4.5217	3.0436	0.93	0.3388	0.0780
142 * <u>Scarus coeruleinus</u>	Midnight parrotfish	1	TL	540	540	-4.8764	3.0618	1.00	0.0000	0.0000
143 <u>Scarus coeruleus</u>	Blue parrotfish	15	TL	103	610	-5.0162	3.1109	0.99	0.0838	0.0558
144 <u>Scarus croicensis</u>	Striped parrotfish	7	TL	24	96	-4.8887	3.0548	1.00	0.0628	0.0341

Table 1. (continued) - Summary of southern Florida reef fish weight-regressions for 7,503 fish.

Regression formula: $\log \text{WEIGHT(gms)} = \log a + b \log \text{LENGTH(mm)}$.

Names of fishes are from Robins et. al. (1980).

* - Species with less than four actual fish measurements. The regression values presented can be only considered to represent very crude estimates. The regression lines were derived by forcing the intercept through a point at or close to the origin by adding a "zero" data point.

SPECIES	COMMON NAME	SIZE (mm)					STANDARD DEVIATION			
		NO.	TYPE	MIN.	MAX.	$\log a$	b	R^2	b	$\log a$
145 <i>Scarus guacamaia</i>	Rainbow parrotfish	11	TL	128	484	-4.8714	3.0626	0.99	0.1026	0.0553
146 <i>Scarus taeniopterus</i>	Princess parrotfish	4	TL	176	280	-4.1836	2.7086	0.85	0.8116	0.1289
147 * <i>Scomberomorus cavalla</i>	King mackerel	1	FL	510	510	-5.0538	2.9731	1.00	0.0000	0.0000
148 * <i>Scomberomorus maculatus</i>	Spanish mackerel	1	FL	472	472	-5.0356	2.9822	1.00	0.0000	0.0000
149 <i>Scorpaena plumieri</i>	Scorpion fish	122	TL	13	340	-4.5626	2.9486	0.99	0.0323	0.0582
150 <i>Selene vomer</i>	Lookdown	31	FL	127	295	-4.7842	3.0758	0.97	0.1039	0.0409
151 <i>Seriola dumerili</i>	Greater amberjack	30	FL	220	728	-4.2985	2.8091	0.98	0.0693	0.0389
152 * <i>Seriola rivoliana</i>	Almaco jack	2	FL	258	307	-4.8908	3.0548	1.00	0.0100	0.0364
153 * <i>Serranus baldwini</i>	Lanternfish	1	FL	51	51	-4.9285	3.0358	1.00	0.0000	0.0000
154 <i>Serranus tigrinus</i>	Harlequin bass	7	FL	30	288	-4.8862	3.0475	1.00	0.0529	0.0475
155 * <i>Sparisoma atomarium</i>	Greenblotch parrotfish	2	TL	62	86	-4.9446	3.0275	1.00	0.0041	0.0131
156 <i>Sparisoma aurofrenatum</i>	Redband parrotfish	17	FL	129	235	-5.7587	3.4291	0.88	0.3336	0.0948
157 <i>Sparisoma chrysopterum</i>	Redtail parrotfish	228	FL	27	395	-5.1754	3.1708	0.98	0.0321	0.0704
158 * <i>Sparisoma rubripinne</i>	Yellowtail parrotfish	2	FL	240	315	-4.8701	3.0641	1.00	0.0241	0.0872
159 <i>Sparisoma viride</i>	Stoplight parrotfish	67	FL	129	382	-4.5223	2.9214	0.96	0.0707	0.0370
160 <i>Sphoeroides nephelus</i>	Southern puffer	23	TL	155	255	-4.0772	2.6973	0.72	0.3711	0.1228
161 <i>Sphoeroides spengleri</i>	Bandtail puffer	40	TL	27	303	-5.2360	3.2671	0.96	0.1065	0.1598
162 * <i>Sphoeroides testudineus</i>	Checkered puffer	3	TL	170	205	-4.8561	3.0717	1.00	0.0083	0.0305
163 <i>Sphyraena barracuda</i>	Barracuda	10	FL	58	1020	-5.3865	3.0825	0.99	0.0850	0.1051
164 * <i>Sphyraena picudilla</i>	Southern sennet	1	FL	430	430	-5.1165	2.9417	1.00	0.0000	0.0000
165 <i>Starksia ocellata</i>	Checkered blenny	19	TL	10	45	-4.7368	2.8771	0.69	0.4653	0.3248
166 * <i>Synodus foetens</i>	Inshore lizardfish	1	FL	305	305	-5.0425	2.9787	1.00	0.0000	0.0000
167 * <i>Synodus intermedius</i>	Sand diver	3	FL	299	386	-5.0020	2.9988	1.00	0.0042	0.0165
168 <i>Thalassoma bifasciatum</i>	Bluehead	39	TL	15	118	-4.8865	2.9162	0.90	0.1588	0.2709
169 * <i>Umbrina coroides</i>	Sand drum	1	TL	192	192	-4.9506	3.0247	1.00	0.0000	0.0000
170 <i>Urolophus jamaicensis</i>	Yellow stingray	6	TL	273	416	-5.2244	3.0826	0.99	0.1563	0.0227
171 * <i>Vomer setapinnis</i>	Atlantic moonfish	2	FL	187	224	-4.8312	3.0856	1.00	0.0245	0.0852
172 * <i>Xyrichtys martinicensis</i>	Rosy razorfish	3	TL	25	62	-4.8221	3.0780	1.00	0.0593	0.1879
173 <i>Xyrichtys novacula</i>	Pearly razorfish	5	TL	35	100	-3.5613	2.2430	0.97	0.2352	0.0792
174 * <i>Xyrichtys splendens</i>	Green razorfish	3	TL	97	104	-5.0012	2.9995	1.00	0.0121	0.0420

Table 2. - Summary of Caribbean (Puerto Rico, St. Thomas/St. John, and St.Croix) reef fish weight-length regressions for 24,383 fish. Regression formula: $\log W(\text{gms}) = \log a + b \log L(\text{mm})$.

SPECIES Common Name	AREA	NO.	TYPE	SIZE (mm)			log a	b	R^2
				MIN.	MAX.				
1 <i>Acanthurus chirurgus</i> Doctorfish	St. Thomas/St. John	139	FL	160	320	-4.0350	2.7443	0.89	
	St. Croix	227	FL	175	337	-2.4262	2.0768	0.75	
2 <i>Acanthurus coeruleus</i> Blue tang	St. Thomas/St. John	410	FL	135	285	-3.1415	2.3964	0.80	
	St. Croix	2767	FL	100	295	-2.8242	2.2638	0.61	
3 <i>Balistes vetula</i> Queen triggerfish	Puerto Rico	339	FL	175	546	-4.1822	2.8293	0.91	
	St. Thomas/St. John	509	FL	170	435	-4.0637	2.7837	0.94	
4 <i>Calamus bairdii</i> Jolthead porgy	Puerto Rico	114	FL	170	298	-4.1724	2.8216	0.94	
5 <i>Calamus penna</i> Sheepshead porgy	Puerto Rico	596	FL	134	535	-3.5518	2.5420	0.85	
6 <i>Canthidermis sufflamen</i> Ocean triggerfish	St. Thomas/St. John	3	FL	330	350	-9.0383	4.7279	0.74	
7 <i>Caranx ruber</i> Bar jack	St. Thomas/St. John	32	FL	200	730	-4.6704	2.9545	0.98	
	St. Croix	98	FL	175	380	-4.1896	2.7480	0.90	
8 <i>Epinephelus fulvus</i> Coney	Puerto Rico	583	FL	157	639	-4.6508	2.9330	0.83	
	St. Thomas/St. John	189	FL	190	370	-3.3353	2.4000	0.54	
	St. Croix	1644	FL	105	315	-3.6232	2.5105	0.70	
9 <i>Epinephelus guttatus</i> Red hind	Puerto Rico	723	FL	156	474	-4.4431	2.8386	0.88	
	St. Thomas/St. John	448	FL	205	545	-5.0756	3.1001	0.91	
	St. Croix	567	FL	208	500	-5.3835	3.2304	0.92	
10 <i>Epinephelus striatus</i> Nassau grouper	Puerto Rico	60	FL	210	645	-4.8980	3.0386	0.90	
	St. Thomas/St. John	73	FL	330	770	-5.8455	3.3804	0.60	
	St. Croix	4	FL	320	470	-5.5063	3.2832	0.96	
11 <i>Etelis oculatus</i> Queen snapper	St. Thomas/St. John	21	FL	360	890	-3.6326	2.5498	0.85	
	St. Croix	48	FL	200	702	-3.7621	2.5775	0.95	
12 <i>Haemulon flavolineatum</i> French grunt	Puerto Rico	200	FL	137	296	-3.8605	2.6509	0.76	
	St. Thomas/St. John	12	FL	170	210	-3.4610	2.4850	0.72	
	St. Croix	232	FL	170	280	-3.5676	2.5255	0.54	
13 <i>Haemulon plumieri</i> White grunt	Puerto Rico	1071	FL	135	444	-4.2654	2.8186	0.89	
	St. Thomas/St. John	39	FL	190	700	-4.7822	3.0336	0.97	
	St. Croix	1588	FL	50	325	-2.2775	1.9849	0.57	
14 <i>Haemulon sciurus</i> Bluestriped grunt	St. Thomas/St. John	23	FL	205	310	-4.6001	2.9630	0.91	
	St. Croix	138	FL	193	285	-3.9303	2.6930	0.68	
15 <i>Holocentrus ascensionis</i> Squirrelfish	St. Thomas/St. John	98	FL	180	270	-3.3204	2.4280	0.66	
	St. Croix	187	FL	160	250	-2.6159	2.1061	0.38	
16 <i>Lachnolaimus maximus</i> Hogfish	Puerto Rico	53	FL	230	595	-3.9814	2.7055	0.81	
	St. Thomas/St. John	27	FL	245	704	-4.3997	2.8828	0.99	
	St. Croix	3	FL	308	363	-5.3159	3.2641	1.00	
17 <i>Lactophrys polygona</i> Honeycomb cowfish	Puerto Rico	203	FL	132	417	-3.9652	2.6794	0.91	
	St. Croix	199	FL	135	385	-3.4940	2.4922	0.86	
	Puerto Rico	219	FL	120	360	-2.9762	2.2372	0.81	
18 <i>Lactophrys quadricornis</i> Scrawled cowfish	St. Croix	9	FL	244	330	-5.8621	3.4177	0.91	

Table 2. (continued) - Summary of Caribbean (Puerto Rico, St. Thomas/St. John, and St.Croix) reef fish weight-length regressions for 24,383 fish. Regression formula: $\log W(\text{gms}) = \log a + b \log L(\text{mm})$.

SPECIES Common Name	AREA	NO.	TYPE	SIZE (mm)		$\log a$	b	R^2
				MIN.	MAX.			
19 <u>Lutjanus analis</u> Mutton snapper	Puerto Rico	53	FL	214	771	-4.6555	2.9502	0.94
	St. Thomas/St. John	27	FL	230	610	-5.2541	3.1747	0.99
	St. Croix	17	FL	260	630	-4.8341	3.0345	0.96
20 <u>Lutjanus apodus</u> Schoolmaster	Puerto Rico	72	FL	185	464	-5.1010	3.1541	0.98
	St. Thomas/St. John	39	FL	255	450	-4.7653	3.0096	0.94
	St. Croix	101	FL	188	470	-4.2103	2.7965	0.81
21 <u>Lutjanus buccanella</u> Blackfin snapper	Puerto Rico	20	FL	173	360	-4.4161	2.8593	0.98
	St. Thomas/St. John	180	FL	190	450	-4.1265	2.7346	0.91
	St. Croix	65	FL	212	440	-4.6475	2.9636	0.95
22 <u>Lutjanus synagris</u> Lane snapper	Puerto Rico	396	FL	156	414	-4.4119	2.8437	0.85
	St. Thomas/St. John	99	FL	175	400	-4.7458	2.9809	0.94
	St. Croix	4	FL	185	238	-2.3588	1.9954	0.67
23 <u>Lutjanus vivanus</u> Silk snapper	Puerto Rico	181	FL	149	403	-5.3646	3.2368	0.93
	St. Thomas/St. John	36	FL	195	635	-4.6001	2.9132	0.97
	St. Croix	165	FL	222	650	-4.2096	2.7812	0.93
24 <u>Mulloidichthys martinicus</u> Yellow goatfish	Puerto Rico	90	FL	164	290	-4.9593	3.0918	0.93
	St. Thomas/St. John	22	FL	225	325	-2.6201	2.1091	0.64
	St. Croix	547	FL	125	270	-3.2528	2.3772	0.55
25 <u>Mycteroperca venenosa</u> Yellowfin grouper	Puerto Rico	19	FL	225	550	-4.7033	2.9764	0.93
	St. Thomas/St. John	103	FL	285	900	-5.1611	3.1402	0.96
	St. Croix	11	FL	254	645	-3.2945	2.3892	0.75
26 <u>Ocyurus chrysurus</u> Yellowtail snapper	Puerto Rico	992	FL	29	562	-3.1552	2.3311	0.76
	St. Thomas/St. John	456	FL	215	560	-4.1770	2.7588	0.92
	St. Croix	610	FL	194	562	-4.3379	2.8110	0.96
27 <u>Pomacanthus arcuatus</u> Gray angelfish	St. Thomas/St. John	82	FL	25	490	0.8074	0.8716	0.26
28 <u>Pseudupeneus maculatus</u> Spotted goatfish	Puerto Rico	1160	FL	127	297	-4.6397	2.9581	0.85
	St. Thomas/St. John	3	FL	195	270	-6.7109	3.8059	1.00
	St. Croix	125	FL	180	262	-3.7431	2.5866	0.74
29 <u>Rhomboplites aurorubens</u> Vermilion snapper	Puerto Rico	162	FL	152	303	-4.6686	2.9512	0.87
	St. Thomas/St. John	6	FL	150	245	-5.0857	3.1214	0.91
	St. Croix	14	FL	310	405	-3.5651	2.5201	0.95
30 <u>Scarus taeniopterus</u> Princess parrotfish	St. Thomas/St. John	29	FL	215	300	-3.2257	2.3852	0.87
	St. Croix	334	FL	200	350	-2.0098	1.8751	0.43
31 <u>Sparisoma aurofrenatum</u> Redband parrotfish	St. Thomas/St. John	25	FL	180	330	-3.1111	2.3363	0.84
	St. Croix	434	FL	190	280	-4.0781	2.7438	0.43
32 <u>Sparisoma chrysopterum</u> Redtail parrotfish	St. Thomas/St. John	93	FL	220	425	-3.3969	2.4533	0.66
	St. Croix	1970	FL	150	383	-3.0509	2.3192	0.58
32 <u>Sparisoma viride</u> Stoplight parrotfish	St. Thomas/St. John	53	FL	218	460	-5.0057	3.1207	0.93
	St. Croix	1693	FL	190	425	-4.4317	2.9051	0.84

Table 3. - List of species observed in visual censusing by Reef Team and for which length-weight data is needed (fewer than 30 measured individuals available).

SPECIES	COMMON NAME
<u>Acanthemblemaria aspera</u>	Roughhead blenny
<u>Acanthemblemaria chaplini</u>	Papillose blenny
<u>Acanthurus bahianus</u>	Ocean surgeon
<u>Adioryx vexillarius</u>	Dusky squirrelfish
<u>Aetobatus narinari</u>	Spotted eagle ray
<u>Alectis crinitus</u>	African pompano
<u>Amblycirrhitus pinos</u>	Redspotted hawkfish
<u>Anchoa lyolepis</u>	Dusky anchovy
<u>Anisotremus surinamensis</u>	Black margate
<u>Apogon binotatus</u>	Barred cardinalfish
<u>Apogon maculatus</u>	Flamefish
<u>Apogon quadrisquamatus</u>	Sawcheek cardinalfish
<u>Archosargus probatocephalus</u>	Sheepshead
<u>Archosargus rhomboidalis</u>	Sea bream
<u>Astrapogon stellatus</u>	Conchfish
<u>Atherinomorus stipes</u>	Hardhead silverside
<u>Aulostomus maculatus</u>	Trumpetfish
<u>Blennius cristata</u>	Molly miller
<u>Bodianus puchellus</u>	Spotfin hogfish
<u>Bodianus rufus</u>	Spanish hogfish
<u>Bothus lunatus</u>	Peacock flounder
<u>Callionymus bairdi</u>	Lancer dragonet
<u>Cantherhines macrocerus</u>	Whitespotted filefish
<u>Cantherhines pullus</u>	Orangespotted filefish
<u>Canthigaster rostrata</u>	Sharpnose puffer
<u>Caranx hippos</u>	Crevalle jack
<u>Caranx latus</u>	Horse-eye jack
<u>Centropomus undecimalis</u>	Snook
<u>Centropristes oxyurus</u>	Bank sea bass
<u>Centropristes striata</u>	Black sea bass
<u>Chaetodon capistratus</u>	Foureye butterflyfish
<u>Chaetodon striatus</u>	Banded butterflyfish
<u>Chilomycterus schoepfi</u>	Stripped burrfish
<u>Chromis cyaneus</u>	Blue chromis
<u>Chromis insolatus</u>	Sunshinefish
<u>Chromis multilineatus</u>	Brown chromis
<u>Chromis scotti</u>	Purple reeffish
<u>Clepticus parrai</u>	Creole wrasse
<u>Coryphopterus dircrus</u>	Colon goby
<u>Coryphopterus glaucofraenum</u>	Bridled goby
<u>Coryphopterus personatus</u>	Masked goby
<u>Cryptotomus roseus</u>	Bluelip parrotfish
<u>Dactylopterus volitans</u>	Flying gurnard
<u>Dasyatis americana</u>	Southern stingray
<u>Decapterus macarellus</u>	Mackerel scad
<u>Decapterus punctatus</u>	Round scad
<u>Echeneis naucrates</u>	Sharksucker

Table 3. (continued) - List of species observed in visual censusing by Reef Team and for which length-weight data is needed (fewer than 30 measured individuals available).

SPECIES	COMMON NAME
<u>Elops saurus</u>	Ladyfish
<u>Emblemaria pandionis</u>	Sailfin blenny
<u>Enchelycore nigrans</u>	Viper moray
<u>Enneanectes boehlkei</u>	Roughhead triplefin
<u>Enneanectes pectoralis</u>	Redeye triplefin
<u>Epinephelus adscensionis</u>	Rock hind
<u>Epinephelus cruentatus</u>	Graysby
<u>Epinephelus itajara</u>	Jewfish
<u>Epinephelus niveatus</u>	Snowy grouper
<u>Epinephelus striatus</u>	Nassau grouper
<u>Equetus acuminatus</u>	High-hat
<u>Equetus lanceolatus</u>	Jackknife-fish
<u>Equetus punctatus</u>	Spotted drum
<u>Eucinostomus argenteus</u>	Spotfin mojarra
<u>Eucinostomus havana</u>	Bigeye mojarra
<u>Gerres cinereus</u>	Yellowfin mojarra
<u>Ginglymostoma cirratum</u>	Nurse shark
<u>Gnatholepis thompsoni</u>	Goldspot goby
<u>Gobiesox strumosus</u>	Skilletfish
<u>Gobiosoma evelynae</u>	Sharpnose goby
<u>Gobiosoma macrodon</u>	Tiger goby
<u>Gobiosoma oceanops</u>	Neon goby
<u>Gramma loreto</u>	Fairy basslet
<u>Gymnothorax funebris</u>	Green moray
<u>Gymnothorax moringa</u>	Spotted moray
<u>Gymnothorax nigromarginatus</u>	Blackedge moray
<u>Gymnothorax saxicola</u>	Ocellated moray
<u>Gymnothorax vicinus</u>	Purplemouth moray
<u>Haemulon album</u>	Margate
<u>Haemulon carbonarium</u>	Caesar grunt
<u>Haemulon chrysargyreum</u>	Smallmouth grunt
<u>Haemulon macrostomum</u>	Spanish grunt
<u>Haemulon striatum</u>	Striped grunt
<u>Halichoeres bivittatus</u>	Slippery dick
<u>Halichoeres garnoti</u>	Yellowhead wrasse
<u>Halichoeres maculipinna</u>	Clown wrasse
<u>Halichoeres pictus</u>	Rainbow wrasse
<u>Halichoeres poeyi</u>	Blackear wrasse
<u>Halichoeres radiatus</u>	Puddingwife
<u>Hemiemblemaria simulus</u>	Wrasse blenny
<u>Hemiramphus brasiliensis</u>	Ballyhoo
<u>Holacanthus tricolor</u>	Rock beauty
<u>Holocentrus coruscus</u>	Reef squirrelfish

Table 3. (continued) - List of species observed in visual censusing by Reef Team and for which length-weight data is needed (fewer than 30 measured individuals available).

SPECIES	COMMON NAME
<u>Holocentrus Marianus</u>	Longjaw squirrelfish
<u>Holocentrus rufus</u>	Longspine squirrelfish
<u>Holocentrus vexillarius</u>	Dusky squirrelfish
<u>Hypseurochilus bermudensis</u>	Barred blenny
<u>Hypoplectrus unicolor</u>	Butter hamlet
<u>Inermia vittata</u>	Boga
<u>Ioglossus calliurus</u>	Blue goby
<u>Jenkinsia lamprotaenia</u>	Dwarf herring
<u>Kyphosus sectatrix</u>	Bermuda chub
<u>Lactophrys trigonus</u>	Trunkfish
<u>Lagodon rhomboides</u>	Pinfish
<u>Liopropoma rubre</u>	Peppermint bass
<u>Lutjanus cyanopterus</u>	Cubera snapper
<u>Lutjanus jocu</u>	Dog snapper
<u>Lutjanus mahogoni</u>	Mahogonay snapper
<u>Lythrypnus spilus</u>	Bluegold goby
<u>Malacanthus plumieri</u>	Sand tilefish
<u>Malacoctenus aurolineatus</u>	Goldline blenny
<u>Malacoctenus gilli</u>	Dusky blenny
<u>Malacoctenus macrops</u>	Rosy blenny
<u>Malacoctenus sp.</u>	Unidentified blenny
<u>Malacoctenus triangulatus</u>	Saddled blenny
<u>Malacoctenus versicolor</u>	Barfin blenny
<u>Manta birostris</u>	Atlantic manta
<u>Megalops atlanticus</u>	Tarpon
<u>Melichthys niger</u>	Black durgon
<u>Microgobius carri</u>	Seminole goby
<u>Microgobius microlepis</u>	Banner goby
<u>Microspathodon chrysurus</u>	Yellowtail damselfish
<u>Monacanthus ciliatus</u>	Fringed filefish
<u>Monacanthus tuckeri</u>	Slender filefish
<u>Mulloidichthys martinicus</u>	Yellow goatfish
<u>Muraena miliaris</u>	Goldentail moray
<u>Muraena retifera</u>	Reticulate moray
<u>Mycteroperca bonaci</u>	Black grouper
<u>Mycteroperca phenax</u>	Scamp
<u>Mycteroperca tigris</u>	Tiger grouper
<u>Mycteroperca venenosa</u>	Yellowfin grouper
<u>Myripristis jacobus</u>	Blackbar soldierfish
<u>Odontoscion dentex</u>	Reef crocker
<u>Ogcocephalus radiatus</u>	Polka-dot batfish







Table 3. (continued) - List of species observed in visual censusing by Reef Team and for which length-weight data is needed (fewer than 30 measured individuals available).

SPECIES	COMMON NAME
<u>Ophioblennius atlanticus</u>	Redlip blenny
<u>Opistognathus aurifrons</u>	Yellowhead jawfish
<u>Opistognathus maxillosus</u>	Mottled jawfish
<u>Opistognathus whitehursti</u>	Dusky jawfish
<u>Paranthias furcifer</u>	Creole-fish
<u>Pempheris schomburgkii</u>	Glassy sweeper
<u>Phaeoptyx pigmentaria</u>	Dusky cardinalfish
<u>Pomacentrus diencaeus</u>	Longfin damselfish
<u>Pomacentrus fuscus</u>	Dusky damselfish
<u>Pomacentrus leucostictus</u>	Beaugregory
<u>Pomacentrus planifrons</u>	Three spot damselfish
<u>Pomacentrus variabilis</u>	Cocoa damselfish
<u>Priacanthus cruentatus</u>	Glasseseye snapper
<u>Prionotus roseus</u>	Bluespotted searobin
<u>Quisquilius hipoliti</u>	Rusty goby
<u>Rachycentron canadum</u>	Cobia
<u>Rypticus saponaceus</u>	Greater soapfish
<u>Scarus coelestinus</u>	Midnight parrotfish
<u>Scarus coeruleus</u>	Blue parrotfish
<u>Scarus croicensis</u>	Striped parrotfish
<u>Scarus guacamaia</u>	Rainbow parrotfish
<u>Scarus taeniopterus</u>	Princess parrotfish
<u>Scarus vetula</u>	Queen parrotfish
<u>Scomberomorus cavalla</u>	King mackerel
<u>Scomberomorus maculatus</u>	Spanish mackerel
<u>Scomberomorus regalis</u>	Cero mackerel
<u>Seriola rivoliana</u>	Almaco jack
<u>Serranus baldwini</u>	Lanternfish
<u>Serranus tabacarius</u>	Tobaccofish
<u>Serranus tigrinus</u>	Harlequin bass
<u>Serranus tortugaram</u>	Chalk bass
<u>Sparisoma atomarium</u>	Greenblotch parrotfish
<u>Sparisoma radians</u>	Bucktooth parrotfish
<u>Sparisoma rubripinne</u>	Yellowtail parrotfish
<u>Sphoeroides nephelus</u>	Southern puffer
<u>Sphoeroides testudineus</u>	Checkered puffer
<u>Sphyraena barracuda</u>	Barracuda
<u>Sphyraena picudilla</u>	Southern sennet
<u>Sphyraena mokarran</u>	Great hammerhead
<u>Starksia ocellata</u>	Checkered blenny
<u>Strongylura notata</u>	Redfin needlefish
<u>Strongylura timucu</u>	Timucu
<u>Synodus foetens</u>	Inshore lizardfish
<u>Synodus intermedius</u>	Sand diver
<u>Trachinotus falcatus</u>	Permit







Table 3. (continued) - List of species observed in visual censusing by Reef Team and for which length-weight data is needed (fewer than 30 measured individuals available).

SPECIES	COMMON NAME
<u>Tylosurus crocodilus</u>	Houndfish
<u>Umbrina coroides</u>	Sand drum
<u>Urolophus jamaicensis</u>	Yellow stingray
<u>Vomer setapinnis</u>	Atlantic moonfish
<u>Xyrichtys martinicensis</u>	Rosy razorfish
<u>Xyrichtys novacula</u>	Pearly razorfish
<u>Xyrichtys splendens</u>	Green razorfish







Appendix A.- Frequency distribution and regression graphics for southern Florida fish species with >30 measurements.









Appendix A.— Frequency distribution and regression graphics for
 (cont.) southern Florida fish species with >30 measurements.









Appendix A.- Frequency distribution and regression graphics for
 (cont.) southern Florida fish species with >30 measurements.









Appendix A.— Frequency distribution and regression graphics for
 (cont.) southern Florida fish species with >30 measurements.









Appendix A.- Frequency distribution and regression graphics for
 (cont.) southern Florida fish species with >30 measurements.









Appendix A.- Frequency distribution and regression graphics for
 (cont.) southern Florida fish species with >30 measurements.









Appendix A.- Frequency distribution and regression graphics for
 (cont.) southern Florida fish species with >30 measurements.









Appendix A.— Frequency distribution and regression graphics for
 (cont.) southern Florida fish species with >30 measurements.









Appendix A.— Frequency distribution and regression graphics for
 (cont.) southern Florida fish species with >30 measurements.









Appendix A.- Frequency distribution and regression graphics for
 (cont.) southern Florida fish species with >30 measurements.









Appendix A.- Frequency distribution and regression graphics for
 (cont.) southern Florida fish species with >30 measurements.









Appendix A.- Frequency distribution and regression graphics for
 (cont.) southern Florida fish species with >30 measurements.









Appendix A.— Frequency distribution and regression graphics for
 (cont.) southern Florida fish species with >30 measurements.









Appendix A.— Frequency distribution and regression graphics for
 (cont.) southern Florida fish species with >30 measurements.



Appendix A.— Frequency distribution and regression graphics for
 (cont.) southern Florida fish species with >30 measurements.



Appendix A.— Frequency distribution and regression graphics for
 (cont.) southern Florida fish species with >30 measurements.



Appendix A.- Frequency distribution and regression graphics for
 (cont.) southern Florida fish species with >30 measurements.

